

# NAUB JOURNAL OF SCIENCE AND TECHNOLOGY

(NAUBJOST)

e-ISSN:2811-2350



Email: [jost@naub.edu.ng](mailto:jost@naub.edu.ng)

<http://journal.naub.edu.ng/naubjournal/>

## **MODIFICATION OF THE INTERIOR SOLUTION OF EINSTEIN'S G22 FIELD EQUATION FOR A HOMOGENEOUS SPHERICAL MASSIVE BODIES WHOSE FIELDS DIFFER IN RADIAL SIZE, POLAR ANGLE AND TIME**

<sup>1</sup>Rilwan Usman., <sup>2</sup>A. U. Maisalatee, <sup>3</sup>M.M. Idris, <sup>4</sup>A.A. Bello, <sup>5</sup>A. Ubaidullah and <sup>3</sup>O.G. Okara

<sup>1</sup>Department of Physics, Nigerian Army University, PMB 1500 Biu, Borno State, Nigeria.

<sup>2</sup>Liyu Unity Science Academy, Behind Yaro Sule Filling Station, P.M.B 03 Keffi, Nasarawa State, Nigeria.

<sup>3</sup>Department of Physics, Nasarawa State University, Keffi, P.M.B. 1022, Nigeria.

<sup>4</sup>Department of Physics, Federal University, Lafia, Nasarawa State, Nigeria.

<sup>5</sup>Department of Physics, Federal University Dutsin-ma, P.M.B 5001 Dutsin-ma, Katsina, Nigeria.

E-mail: [rilwan.usman@naub.edu.ng](mailto:rilwan.usman@naub.edu.ng)

Phone Number: 08037772627/07055324002

### **Abstract**

In general theory of relativity, Einstein's field equations relate the geometry of space-time with the distribution of matter within it. Research has showed that the tensors for spherical massive bodies are not functions of radial distance only as shown by Schwarzschild; they depend on other factors such as polar angle, azimuthal angle and time. In this article, the analytical interior solution of Einstein's field equation for homogeneous spherical body whose tensor field varies with time, radial distance and polar angle using weak field and slow-motion approximation was formulated. The obtained result converges to Newton's dynamical scalar potential with additional time factor not found in the well-known Newton's dynamical theory of gravitation which is a profound discovery with the dependency on three arbitrary functions. The result obtained can be used in the study of rotating astrophysical bodies such as stars. The result obeyed the equivalence principle of Physics.

**Keywords:** Schwarzschild metric, Einstein Equation, radial size, polar angle, Einstein tensor

### **Introduction**

According to Einstein's theory of gravitation published in 1915/1916, which unifies Special Relativity and Sir Isaac Newton's law of universal gravitation, that gravitation is not due to a force but rather a manifestation of curved space and time, with the curvature being produced by the mass-energy and momentum content of the space-time. This is termed General Relativity and is the most widely accepted theory of gravitation (Howusu, 2010; Chifu, 2012). The equations are in the form of tensor equation which related the local space-time curvature (expressed by the Einstein tensor) with the local energy and momentum within that space-time (expressed by the stress-energy tensor) (Misner *et al.*, 1973). Einstein's General Relativity is the leading theory of space-time and gravity. Exact solutions of Einstein's equations thus model gravitating systems and enable exploration of the Mathematics and Physics of the theory (Malcolm, 2003). After the publication of Einstein's geometrical

gravitational field equations in 1915, the search for their exact and analytical solutions for all the gravitational fields in nature began (Chifu and Howusu 2009; Howusu, 2010; Chifu, 2012). Schwarzschild first constructed the exact solution to this field equation in static and pure radial spherical polar coordinates in 1916 by considering astrophysical bodies such as the sun and the stars (Lumbi *et al.*, 2014). In Schwarzschild's metric, the tensor field varies with radial distance only.

A new method and approach was introduced to formulate exact analytical solutions (Chifu, 2009) as an extension of Schwarzschild's method. This new approach took into consideration the fact that tensor field of astrophysical bodies does not depend on radial distance only as indicated in Schwarzschild's equation. This new approach was used in several studies of Einstein's geometrical field equations such as (Chifu and Lumbi, 2008; Chifu and Howusu 2009; Sarki *et al.*, 2018). This method would help in the study of Ceres, Pluto, Makemake, Haumea, the Oort Cloud and other astrophysical bodies. In this research work, we show how exact analytical solution of the interior field equation can be constructed in the limit of  $c^{-2}$  in a gravitational field for time varying spherical massive bodies using the new method and approach.

### Einstein's Equation

Einstein's field equation (EFE) interior to a homogeneous spherical distribution of mass is given generally as (Misner *et al.*, 1973; Tajmar, 2001; Howusu, 2008; Chifu, 2009)

$$R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} = \frac{8\pi GT_{\mu\nu}}{c^4} \quad (1)$$

Where  $T_{\mu\nu}$  is the energy-momentum tensor due to any distribution of mass or pressure

G is the universal gravitational constant.

Now, let us assume that the homogeneous mass distribution is in a weak field limit. We can neglect the contribution from the source, thus define the energy-momentum tensor given as

$$T_{\mu\nu} = \frac{1}{2}\rho_o c^2 \quad (2)$$

$$G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} = \frac{4\pi G\rho_o}{c^2} \quad (3)$$

Where  $\rho_o$  is the density

$c$  is the speed of light in vacuum.

It was observed in (Misner *et al.*, 1973) that the exterior field equations along the  $G_{00}$ ,  $G_{22}$  and  $G_{33}$  converge within the exterior field, similarly along the interior field.

For mathematical convenience, we choose  $G_{22}$

Hence the non-trivial field equation is

$$R_{22} - \frac{1}{2}Rg_{22} = \frac{4\pi G\rho_o}{c^2} \quad (4)$$

The affine connections of this field constructed by (Lumbi *et al.*, 2014) were used to construct the Ricci tensor  $R_{22}$  and the curvature scalar  $R$  given respectively as:

$$R_{22} = R_{220}^0 + R_{221}^1 + R_{222}^2 + R_{223}^3 \quad (5)$$

$$R = g^{00}R_{00} + g^{11}R_{11} + g^{22}R_{22} + g^{33}R_{33} \quad (6)$$

The expanded form of equations (5) and (6) are given as

$$\begin{aligned} R_{22} = & \Gamma_{20,2}^0 + \Gamma_{20}^0\Gamma_{02}^0 - \Gamma_{22}^1\Gamma_{10}^0 + \Gamma_{21,2}^1 - \Gamma_{22,1}^1 + \Gamma_{21}^1\Gamma_{12}^1 - \Gamma_{22}^1\Gamma_{11}^1 - \Gamma_{12}^2\Gamma_{22}^1 \\ & + \Gamma_{22,2}^2 - \Gamma_{22,2}^2 + \Gamma_{22}^0\Gamma_{02}^2 - \Gamma_{22}^0\Gamma_{02}^2 + \Gamma_{22}^1\Gamma_{12}^2 - \Gamma_{22}^1\Gamma_{12}^2 + \Gamma_{22}^2\Gamma_{22}^2 - \Gamma_{22}^2\Gamma_{22}^2 + \Gamma_{22}^3\Gamma_{32}^2 - \Gamma_{22}^3\Gamma_{32}^2 \\ & + \Gamma_{23,2}^3 - \Gamma_{22}^1\Gamma_{13}^3 + \Gamma_{23}^3\Gamma_{32}^3 \end{aligned} \quad (7)$$

$$\begin{aligned} R = & g^{00} \{ \Gamma_{00,0}^0 - \Gamma_{00,0}^0 + \Gamma_{00}^0\Gamma_{00}^0 - \Gamma_{00}^0\Gamma_{00}^0 + \Gamma_{00}^1\Gamma_{10}^0\Gamma_{01,0}^1 - \Gamma_{00}^1\Gamma_{10}^0 \\ & - \Gamma_{00}^2\Gamma_{20}^0 - \Gamma_{00}^2\Gamma_{20}^0 + \Gamma_{00}^3\Gamma_{30}^0 - \Gamma_{00}^3\Gamma_{30}^0 - \Gamma_{00,1}^1 + \Gamma_{01}^0\Gamma_{10}^0 - \Gamma_{00}^0\Gamma_{01}^1 + \Gamma_{01}^1\Gamma_{10}^0 \\ & - \Gamma_{00}^1\Gamma_{11}^1 - \Gamma_{00}^2\Gamma_{21}^1 - \Gamma_{00,2}^2 + \Gamma_{02}^0\Gamma_{20}^0 - \Gamma_{00}^1\Gamma_{12}^2 - \Gamma_{00}^1\Gamma_{13}^3 - \Gamma_{00}^2\Gamma_{23}^3 \} \\ & + g^{11} \{ \Gamma_{10,1}^0 - \Gamma_{11,0}^0 + \Gamma_{10}^0\Gamma_{01}^0 - \Gamma_{11}^0\Gamma_{00}^0 + \Gamma_{10}^1\Gamma_{11}^0 - \Gamma_{11}^1\Gamma_{10}^0 \\ & - \Gamma_{11}^2\Gamma_{20}^0 + \Gamma_{11,1}^1 - \Gamma_{11,1}^1 + \Gamma_{11}^0\Gamma_{10}^1 - \Gamma_{11}^0\Gamma_{10}^1 + \Gamma_{11}^1\Gamma_{11}^1 - \Gamma_{11}^1\Gamma_{11}^1 + \Gamma_{11}^2\Gamma_{21}^1 - \Gamma_{11}^2\Gamma_{21}^1 \\ & + \Gamma_{11}^3\Gamma_{31}^1 - \Gamma_{11}^3\Gamma_{31}^1 + \Gamma_{12,1}^2 - \Gamma_{11,2}^2 + \Gamma_{12}^1\Gamma_{21}^2 - \Gamma_{11}^1\Gamma_{12}^2 + \Gamma_{12}^2\Gamma_{21}^2 + \Gamma_{13,1}^3 - \Gamma_{11}^1\Gamma_{13}^3 - \Gamma_{11}^2\Gamma_{23}^3 + \Gamma_{13}^3\Gamma_{31}^3 \} \\ & + g^{22} \{ \Gamma_{20,2}^0 + \Gamma_{20}^0\Gamma_{02}^0 - \Gamma_{22}^1\Gamma_{10}^0 + \Gamma_{21,2}^1 - \Gamma_{22,1}^1 + \Gamma_{21}^1\Gamma_{12}^1 - \Gamma_{22}^1\Gamma_{11}^1 - \Gamma_{12}^2\Gamma_{22}^1 \\ & + \Gamma_{22,2}^2 - \Gamma_{22,2}^2 + \Gamma_{22}^0\Gamma_{02}^2 - \Gamma_{22}^0\Gamma_{02}^2 + \Gamma_{22}^1\Gamma_{12}^2 - \Gamma_{22}^1\Gamma_{12}^2 + \Gamma_{22}^2\Gamma_{22}^2 - \Gamma_{22}^2\Gamma_{22}^2 + \Gamma_{22}^3\Gamma_{32}^2 - \Gamma_{22}^3\Gamma_{32}^2 \\ & + \Gamma_{23,2}^3 - \Gamma_{22}^1\Gamma_{13}^3 + \Gamma_{23}^3\Gamma_{32}^3 \} + g^{33} \{ -\Gamma_{00}^1\Gamma_{13}^3 - \Gamma_{00}^2\Gamma_{23}^3 - \Gamma_{33,1}^1 - \Gamma_{33}^1\Gamma_{11}^1 - \Gamma_{33}^2\Gamma_{21}^2 \\ & + \Gamma_{31}^3\Gamma_{13}^1 - \Gamma_{33,2}^2 - \Gamma_{33}^1\Gamma_{12}^2 + \Gamma_{32}^3\Gamma_{23}^3 + \Gamma_{33,3}^3 - \Gamma_{33,3}^3 + \Gamma_{33}^0\Gamma_{03}^3 - \Gamma_{33}^0\Gamma_{03}^3 \\ & + \Gamma_{33}^1\Gamma_{13}^3 - \Gamma_{33}^1\Gamma_{13}^3 + \Gamma_{33}^2\Gamma_{23}^3 - \Gamma_{33}^2\Gamma_{23}^3 + \Gamma_{33}^3\Gamma_{33}^3 - \Gamma_{33}^3\Gamma_{33}^3 \} \end{aligned} \quad (8)$$

Explicitly equations (7) and (8) are given as

$$R_{22} = \frac{2}{c^4} \left( 1 + \frac{2f(t,r,\theta)}{c^2} \right)^{-2} \left( \frac{\partial f(t,r,\theta)}{\partial \theta} \right)^2 + \frac{2r}{c^2} \frac{\partial f(t,r,\theta)}{\partial r} + \frac{2f(t,r,\theta)}{c^2} \quad (9)$$

$$\begin{aligned} R = & \frac{8}{c^4} \left( 1 + \frac{2f(t,r,\theta)}{c^2} \right)^{-3} \left[ \frac{\partial f(t,r,\theta)}{\partial t} \right]^2 - \frac{8}{c^2 r} \frac{\partial f(t,r,\theta)}{\partial r} - \frac{2}{c^2} \left( 1 + \frac{2f(t,r,\theta)}{c^2} \right)^{-2} \left( \frac{\partial^2 f(t,r,\theta)}{\partial t^2} \right) \\ & - \frac{2}{c^2} \left( \frac{\partial^2 f(t,r,\theta)}{\partial r^2} \right) - \frac{2}{c^4 r^2} \left( 1 + \frac{2f(t,r,\theta)}{c^2} \right)^{-2} \left( \frac{\partial f(t,r,\theta)}{\partial \theta} \right)^2 - \frac{4f(t,r,\theta)}{c^2 r^2} \end{aligned} \quad (10)$$

Substituting (9) and (10) into (4) gives

$$\nabla^2 f(t,r,\theta) + \left( \frac{\partial^2 f(t,r,\theta)}{\partial t^2} \right) - \frac{1}{c^2 r^2} \left( \frac{\partial f(t,r,\theta)}{\partial \theta} \right)^2 + \frac{4f(t,r,\theta)}{c^4 r^2} \left( \frac{\partial f(t,r,\theta)}{\partial \theta} \right)^2$$

$$-\frac{4}{c^2} \left[ \frac{\partial f(t,r,\theta)}{\partial t} \right]^2 + \frac{24f(t,r,\theta)}{c^4} \left[ \frac{\partial f(t,r,\theta)}{\partial t} \right]^2 - \frac{4f(t,r,\theta)}{c^2} \left( \frac{\partial^2 f(t,r,\theta)}{\partial t^2} \right) \Bigg] = \frac{4\pi G\rho_o}{c^2} \quad (11)$$

$$\nabla^2 f(t,r,\theta) + \left( \frac{\partial^2 f(t,r,\theta)}{\partial t^2} \right) - \frac{1}{c^2 r^2} \left( \frac{\partial f(t,r,\theta)}{\partial \theta} \right)^2 - \frac{4}{c^2} \left[ \frac{\partial f(t,r,\theta)}{\partial t} \right]^2 - \frac{4f(t,r,\theta)}{c^2} \left( \frac{\partial^2 f(t,r,\theta)}{\partial t^2} \right) \Bigg] = \frac{4\pi G\rho_o}{c^2} \quad (12)$$

To the weak field limit of  $c^0$ , the equation reduces to

$$\nabla^2 f(t,r,\theta) + \frac{\partial^2 f(t,r,\theta)}{\partial t^2} = \frac{4\pi G\rho_o}{c^2} \quad (13)$$

### Conclusion

The result obtained from equation 13, is the Newton dynamical scalar field equation with an additional time factor which signifies the dynamical nature of the system. It is indeed a profound discovery, it confirms the assumption made by (Sarki *et al.*, 2018); that Newton dynamical theory of gravitation (NDTG) is a limiting case of Einstein's geometrical gravitational field equations (EGGFE), and this shades more light on the report of (Kumar *et al.*, 2012). Experimentally established equivalence principle of physics is shown with the dependency of the scalar function on time, radial distance and polar angle.

If the pressure is negligible compared to mass density,

$$\rho_o \equiv 0$$

Our scalar potential will be

$$\nabla^2 f(t,r,\theta) + \frac{\partial^2 f(t,r,\theta)}{\partial t^2} = -\frac{GM_0}{r} \quad (14)$$

Where  $GM_0 = k$

The EFE reduces to Newton's law of gravitation by using both weak field approximation and slow motion approximation, thus equation 3 further splits into four non-linear equations, analogous to Maxwell's equation and could be applied in the study of the Gravitoelectric and Gravitomagnetic coupling phenomena.

## References

- Chifu, E. N. & Lumbi, W. L. (2008). General relativistic equations of motion for test particles exterior to astrophysically real or hypothetically spherical distribution of mass whose tensor field varies with azimuthal angle only. *Continental J. Applied Sciences*, **3**(8):32-38
- Chifu, E. N. (2012). Gravitational Fields Exterior to a Homogeneous Spherical Masses. *The Abraham Zelmanov Journal*, **5**: 31-67
- Chifu, E.N. (2009). Astrophysical Satisfactory Solutions to Einstein  $R_{33}$  Gravitational Field Equations Exterior/Interior to Static Homogeneous Oblate Spheroidal Masses, In Proceedings of 3rd International IMBIC Conference, 75-89.
- Chifu, E.N.& Howusu, S.X.K. (2009). Solution of Einstein's Geometrical Field Equations Exterior to Astrophysical Real or Hypothetical Time-Varying Distributions of Mass within Regions of Spherical Geometry,*Progress in Physics*,**3**:45-48.
- Howusu, S. X. K. (2010). Exact Analytical Solutions of Einstein's Geometrical Gravitational Field Equations, Jos University Press Ltd., 2010, vii-43
- Kumar, K.N.P., Kiranagi, B.S., & Begewadi, C.S. (2012). Einstein Field Equations and Heisenberg's Principle of Uncertainty the Consummation of GTR and Uncertainty Principle, *International Journal of Scientific and Research Publication*, **2**(9):1-56.
- Lumbi, L. W., Ewa, I. I. & Tsaku, N. (2014). Einstein's Equations of Motion for Test Particles Exterior to Spherical Distributions of Mass whose Tensor Field Varies with Time, Radial Distance and Polar Angle. *Archives of Applied Science Research Library*.**6**(5):36-41
- Malcolm, A.H (2013). Exact Solution of Einstein's Equations. *Scholarpedia*, **8**(12):8584
- Misner, C.W., Thorne, K.S., & Wheeler, J.A., Gravitation, 1973, W.H Freeman, San Francisco, USA.
- S.X.K Howusu, Solutions of Einstein's Geometrical Field Equations, 2008, Jos University Press ltd, Plateau State, Nigeria.
- Sarki, M.U., Lumbi, W.L., Ewa, I.I. (2018). Radial Distance and Azimuthal Angle Varying Tensor Field Equation Exterior to a Homogeneous Spherical Mass Distribution, *JNAMP*, **48**: 255-260.
- Tajmar, M. (2001). Coupling of Electromagnetism and Gravitation in the Weak Field Approximation, *Journal of Theoretic*, **3**(1):1-8.